

Notify the Doctor or Go to the Emergency Department if any of the following happens:

- Problems with the incision(s), including increasing pain, swelling, redness, or drainage
- Severe constipation
- Diarrhea (more than three watery stools within 24 hours)
- Difficulty urinating
- Fever greater than 38°C. or 100.4°F.
- Severe bleeding
- Nausea or vomiting
- Chills

After Anal/ Rectal Surgery

Discharge Instructions

For more copies, go online at <http://vch.eduhealth.ca> or email pchem@vch.ca and quote Catalogue No. **ED.150.An13**
© Vancouver Coastal Health, October 2007

The information in this document is intended solely for the person to whom it was given by the health care team.
www.vch.ca

Important Points

Activity

- There is no limit to your activities when you go home.

Baths/Showers

- You may shower and bathe as you like.
- To relieve pain and keep the area clean, sit in a bath of warm water (sometimes called a “sitz bath”) for about 10 minutes 3-4 times a day and after each bowel movement.

Diet

- Remember that a well-balanced diet is important for wound healing.
- Include high fiber and plenty of fluids in your diet to keep the stool soft, especially if taking pain medicine:
 - 6-8 glasses of water daily
 - grain breads
 - bran cereals (All Bran, Bran Buds, Fruit & Fiber)
 - fruits (prunes, raisins, dates, bananas, apples)
 - vegetables

Bowel Function

- Bowel movements after anorectal surgery often cause moderate to severe discomfort. Constipation makes the pain worse and must be avoided. Hard bowel movements may cause trauma to the operative area and slow down healing.
 - **To prevent constipation**, take a bulk-forming stool softener (for example, Metamucil® or Colace). Follow the diet directions (under Diet).
-

Urination

- If you have trouble urinating, do so while sitting in a warm tub of water, or run the water faucet while sitting on the toilet.

Infection/Dressings

- Some bloody discharge after bowel movements is normal for at least two to four weeks after anal/rectal surgery. If you have profuse, continuing bleeding, call your doctor immediately.
- Postoperative infection around the anal opening is not common despite the obvious soiling by stool.
- You can wear a dressing or sanitary napkin to protect your clothing.

Pain

- Use pain medication as ordered by your doctor. This medication may cause constipation.
- Do not drink alcohol when taking pain medication.

Follow-Up Appointment

- Contact your Doctor’s office to arrange for an appointment.
-